

Modelli 1 @ Clamfim

Teoria della misura

Lezione 15

13 ottobre 2014

professor Daniele Ritelli

daniele.ritelli@unibo.it



$$\Gamma\left(\frac{1}{2}\right) = \int_0^{+\infty} t^{\frac{1}{2}-1} e^{-t} dt = \int_0^{\infty} \frac{e^{-t}}{\sqrt{t}} dt$$

$$\Gamma\left(\frac{1}{2}\right) = \int_0^{+\infty} t^{\frac{1}{2}-1} e^{-t} dt = \int_0^{\infty} \frac{e^{-t}}{\sqrt{t}} dt$$

$$\sqrt{t} = u \implies t = u^2 \implies dt = 2u du$$

$$\Gamma\left(\frac{1}{2}\right) = \int_0^{+\infty} t^{\frac{1}{2}-1} e^{-t} dt = \int_0^{\infty} \frac{e^{-t}}{\sqrt{t}} dt$$

$$\sqrt{t} = u \implies t = u^2 \implies dt = 2u du$$

$$\Gamma\left(\frac{1}{2}\right) = \int_0^{\infty} \frac{2ue^{-u^2}}{u} du$$

$$\Gamma\left(\frac{1}{2}\right) = \int_0^{+\infty} t^{\frac{1}{2}-1} e^{-t} dt = \int_0^{\infty} \frac{e^{-t}}{\sqrt{t}} dt$$

$$\sqrt{t} = u \implies t = u^2 \implies dt = 2u du$$

$$\Gamma\left(\frac{1}{2}\right) = \int_0^{\infty} \frac{2ue^{-u^2}}{u} du = 2 \int_0^{\infty} e^{-u^2} du$$

$$\Gamma\left(\frac{1}{2}\right) = \int_0^{+\infty} t^{\frac{1}{2}-1} e^{-t} dt = \int_0^{\infty} \frac{e^{-t}}{\sqrt{t}} dt$$

$$\sqrt{t} = u \implies t = u^2 \implies dt = 2u du$$

$$\Gamma\left(\frac{1}{2}\right) = \int_0^{\infty} \frac{2ue^{-u^2}}{u} du = 2 \int_0^{\infty} e^{-u^2} du = \sqrt{\pi}$$

Una curiosità su Ramanujan e la funzione Gamma

$$\sum_{n=0}^{\infty} (-1)^n \frac{\Gamma^3(n + \frac{1}{2})}{2(n!)^3} (4n + 1) = \sqrt{\pi}$$

Euler Reflexion Formula

$$\Gamma(x)\Gamma(1-x) = \frac{\pi}{\sin \pi x}$$

for $x \notin \mathbb{Z}$

Euler Beta Theorem If $x, y > 0$:

$$\frac{\Gamma(x) \Gamma(y)}{\Gamma(x + y)} = \int_0^1 s^{x-1} (1 - s)^{y-1} ds$$

Euler Beta Theorem If $x, y > 0$:

$$\frac{\Gamma(x) \Gamma(y)}{\Gamma(x+y)} = \int_0^1 s^{x-1} (1-s)^{y-1} ds$$

This identity can be reformulated introducing the Euler Beta function:

$$B(x, y) = \int_0^1 s^{x-1} (1-s)^{y-1} ds$$

in such a way

$$B(x, y) = \frac{\Gamma(x) \Gamma(y)}{\Gamma(x+y)}$$

Proof

We start from Gamma's definition $\Gamma(x) = \int_0^{+\infty} t^{x-1} e^{-t} dt$ then change variable putting $t = u^2$ so that

$$\Gamma(x) = 2 \int_0^{+\infty} u^{2x-1} e^{-u^2} du$$

Proof

We start from Gamma's definition $\Gamma(x) = \int_0^{+\infty} t^{x-1} e^{-t} dt$ then change variable putting $t = u^2$ so that

$$\Gamma(x) = 2 \int_0^{+\infty} u^{2x-1} e^{-u^2} du$$

Similarly

$$\Gamma(y) = 2 \int_0^{+\infty} v^{2y-1} e^{-v^2} dv$$

Proof

We start from Gamma's definition $\Gamma(x) = \int_0^{+\infty} t^{x-1} e^{-t} dt$ then change variable putting $t = u^2$ so that

$$\Gamma(x) = 2 \int_0^{+\infty} u^{2x-1} e^{-u^2} du$$

Similarly

$$\Gamma(y) = 2 \int_0^{+\infty} v^{2y-1} e^{-v^2} dv$$

Now use Fubini's Theorem

$$\Gamma(x)\Gamma(y) = 4 \iint_{[0,+\infty) \times [0,+\infty)} u^{2x-1} v^{2y-1} e^{-(u^2+v^2)} du dv$$

Change to polar coordinate

$$\begin{cases} u = \rho \cos \vartheta \\ v = \rho \sin \vartheta \end{cases}$$

obtaining

Change to polar coordinate

$$\begin{cases} u = \rho \cos \vartheta \\ v = \rho \sin \vartheta \end{cases}$$

obtaining

$$\Gamma(x)\Gamma(y) = 4 \left(\int_0^{+\infty} \rho^{2x+2y-1} e^{-\rho^2} d\rho \right) \left(\int_0^{\pi/2} \cos^{2x-1} \vartheta \sin^{2y-1} \vartheta d\vartheta \right)$$

Change to polar coordinate

$$\begin{cases} u = \rho \cos \vartheta \\ v = \rho \sin \vartheta \end{cases}$$

obtaining

$$\begin{aligned} \Gamma(x)\Gamma(y) &= 4 \left(\int_0^{+\infty} \rho^{2x+2y-1} e^{-\rho^2} d\rho \right) \left(\int_0^{\pi/2} \cos^{2x-1} \vartheta \sin^{2y-1} \vartheta d\vartheta \right) \\ &= \Gamma(x+y) \int_0^{\pi/2} 2 \cos^{2x-1} \vartheta \sin^{2y-1} \vartheta d\vartheta \end{aligned}$$

To end the proof we have to show that

$$B(x, y) = \int_0^{\pi/2} 2 \cos^{2x-1} \vartheta \sin^{2y-1} \vartheta \, d\vartheta$$

To end the proof we have to show that

$$B(x, y) = \int_0^{\pi/2} 2 \cos^{2x-1} \vartheta \sin^{2y-1} \vartheta \, d\vartheta$$

But coming back to Beta's definition

$$B(x, y) = \int_0^1 s^{x-1} (1-s)^{y-1} ds$$

we are done putting $s = \cos^2 \vartheta$

To end the proof we have to show that

$$B(x, y) = \int_0^{\pi/2} 2 \cos^{2x-1} \vartheta \sin^{2y-1} \vartheta \, d\vartheta$$

But coming back to Beta's definition

$$B(x, y) = \int_0^1 s^{x-1} (1-s)^{y-1} ds$$

we are done putting $s = \cos^2 \vartheta$

Conseguenza

$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$

Show, using Gamma and Beta functions, that the Lebesgue measure of $A = \{(x, y) \in \mathbb{R}^2 \mid x^{2/3} + y^{2/3} \leq 1\}$ is $\frac{3}{8}\sqrt{\pi}$

Show, using Gamma and Beta functions, that the Lebesgue measure of $A = \{(x, y) \in \mathbb{R}^2 \mid x^{2/3} + y^{2/3} \leq 1\}$ is $\frac{3}{8}\sqrt{\pi}$




Figure 1: Asteroide

$$m(A) = 4 \int_0^1 \left(1 - x^{2/3}\right)^{3/2} dx$$

$$m(A) = 4 \int_0^1 \left(1 - x^{2/3}\right)^{3/2} dx = 6 \int_0^1 u^{1/2} (1 - u)^{3/2} du$$

$$m(A) = 4 \int_0^1 \left(1 - x^{2/3}\right)^{3/2} dx = 6 \int_0^1 u^{1/2} (1 - u)^{3/2} du = 6B(5/2, 3/2)$$

Evaluate using Beta function

$$\int_0^{\infty} \frac{dx}{1+x^4}$$

Evaluate using Beta function

$$\int_0^{\infty} \frac{dx}{1+x^4}$$

(Smart) Change of variable

$$1+x^4 = \frac{1}{u}$$

Evaluate using Beta function

$$\int_0^{\infty} \frac{dx}{1+x^4}$$

(Smart) Change of variable

$$1+x^4 = \frac{1}{u}$$

$$x = \left(\frac{1-u}{u} \right)^{\frac{1}{4}}$$

Evaluate using Beta function

$$\int_0^{\infty} \frac{dx}{1+x^4}$$

(Smart) Change of variable

$$1+x^4 = \frac{1}{u}$$

$$x = \left(\frac{1-u}{u} \right)^{\frac{1}{4}} \implies dx = -\frac{1}{4} u^{-\frac{5}{4}} (1-u)^{-\frac{3}{4}} du$$

Thus

$$\int_0^\infty \frac{dx}{1+x^4} = \frac{1}{4} \int_0^1 u^{-\frac{1}{4}}(1-u)^{-\frac{3}{4}} du$$

Thus

$$\int_0^{\infty} \frac{dx}{1+x^4} = \frac{1}{4} \int_0^1 u^{-\frac{1}{4}} (1-u)^{-\frac{3}{4}} du$$

Conclusion

$$\int_0^{\infty} \frac{dx}{1+x^4} = \frac{1}{4} B\left(\frac{3}{4}, \frac{1}{4}\right) = \frac{1}{4} \Gamma\left(\frac{3}{4}\right) \Gamma\left(\frac{1}{4}\right)$$

Thus

$$\int_0^{\infty} \frac{dx}{1+x^4} = \frac{1}{4} \int_0^1 u^{-\frac{1}{4}} (1-u)^{-\frac{3}{4}} du$$

Conclusion

$$\int_0^{\infty} \frac{dx}{1+x^4} = \frac{1}{4} B\left(\frac{3}{4}, \frac{1}{4}\right) = \frac{1}{4} \Gamma\left(\frac{3}{4}\right) \Gamma\left(\frac{1}{4}\right)$$

Eventually using the reflexion formula $\Gamma(x)\Gamma(1-x) = \frac{\pi}{\sin \pi x}$ we get

$$\int_0^{\infty} \frac{dx}{1+x^4} = \frac{1}{4} \frac{\pi}{\sin \frac{\pi}{4}} = \frac{\sqrt{2}}{4} \pi$$

Duplication Formula

It is possible to show that

$$2^{2x-1}\Gamma(x)\Gamma(x + \tfrac{1}{2}) = \sqrt{\pi}\Gamma(2x)$$

Tonelli's Theorem Let $A \subset \mathbb{R}^m$ measurable and $f : A \rightarrow \overline{\mathbb{R}}$ measurable. Define \mathcal{S} as the \mathbb{R}^p subset where q -sections of A have positive measure i.e.:

$$\mathcal{S} = \{x \in \mathbb{R}^p \mid \ell_q(A_x) > 0\}$$

Then the equality holds

$$\int_A |f(x, y)| \, dx dy = \int_{\mathcal{S}} \left(\int_{A_x} |f(x, y)| \, dy \right) dx$$

Black Scholes equation:

$$\boxed{\frac{\partial V}{\partial t} + \frac{1}{2}\sigma^2 S^2 \frac{\partial^2 V}{\partial S^2} + rS \frac{\partial V}{\partial S} - rV = 0, \quad S \geq 0, \quad t \in [0, T],} \quad (1)$$

where $V(S, t)$ is the value of the option, S the price of the underlying, t the time, T the expiration date, σ the volatility of the underlying and r the risk-free interest rate. The BS Eq. (1) is a linear parabolic equation of the form

$$\frac{\partial v}{\partial t} = \frac{\partial^2 v}{\partial x^2} + a \frac{\partial v}{\partial x} + bv, \quad (2)$$

which can always be reduced to a diffusion equation

$$\frac{\partial h}{\partial t} = \frac{\partial^2 h}{\partial x^2}. \quad (3)$$

Then an integral form of the solution of the diffusion Eq. (3) can be obtained for any initial condition using for example Fourier transform methods.

Fourier Transform

From: **The Fourier Transform and its Applications** lecture notes
of professor Brad Osgood Stanford University

Let f a real function of a real variable. The Fourier Transform (FT)
of f is the complex valued function:

$$\mathcal{F}f(s) := \int_{-\infty}^{+\infty} e^{-2\pi i s t} f(t) dt$$

Fourier Transform

From: **The Fourier Transform and its Applications** lecture notes
of professor Brad Osgood Stanford University

Let f a real function of a real variable. The Fourier Transform (FT)
of f is the complex valued function:

$$\mathcal{F}f(s) := \int_{-\infty}^{+\infty} e^{-2\pi i s t} f(t) dt$$

There is another notation in use

$$\hat{f}(s) = \int_{-\infty}^{+\infty} e^{-2\pi i s t} f(t) dt$$

A warning on definitions Our definition of the Fourier transform is a standard one, but it's not the only one. The question is where to put the 2π : in the exponential, as we have done; or perhaps as a factor out front; or perhaps left out completely. There's also a question of which is the Fourier transform and which is the inverse, i.e., which gets the minus sign in the exponential.

Summary provided by T. W. Körner Fourier Analysis:

$$\mathcal{F}f(s) = \frac{1}{A} \int_{-\infty}^{+\infty} e^{iBst} f(t) dt$$

Summary provided by T. W. Körner Fourier Analysis:

$$\mathcal{F}f(s) = \frac{1}{A} \int_{-\infty}^{+\infty} e^{iBst} f(t) dt$$

The choices that are found in practice are

$$A = \sqrt{2\pi}$$

$$B = \pm 1$$

$$A = 1$$

$$B = \pm 2\pi$$

$$A = 1$$

$$B = \pm 1$$

Summary provided by T. W. Körner Fourier Analysis:

$$\mathcal{F}f(s) = \frac{1}{A} \int_{-\infty}^{+\infty} e^{iBst} f(t) dt$$

The choices that are found in practice are

$$A = \sqrt{2\pi} \qquad B = \pm 1$$

$$A = 1 \qquad B = \pm 2\pi$$

$$A = 1 \qquad B = \pm 1$$

The definition we've chosen has $A = 1$ and $B = -2\pi$

A sufficient condition for the existence of the integral is $f \in \mathcal{L}^1(\mathbb{R})$. One value is easy to compute, and worth pointing out, namely for $s = 0$ we have

$$\mathcal{F}f(0) := \int_{-\infty}^{+\infty} f(t) dt$$

A sufficient condition for the existence of the integral is $f \in \mathcal{L}^1(\mathbb{R})$. One value is easy to compute, and worth pointing out, namely for $s = 0$ we have

$$\mathcal{F}f(0) := \int_{-\infty}^{+\infty} f(t)dt$$

The inverse Fourier transform is defined by:

$$\mathcal{F}^{-1}g(t) := \int_{-\infty}^{+\infty} e^{2\pi i s t} g(s)ds$$

A sufficient condition for the existence of the integral is $f \in \mathcal{L}^1(\mathbb{R})$. One value is easy to compute, and worth pointing out, namely for $s = 0$ we have

$$\mathcal{F}f(0) := \int_{-\infty}^{+\infty} f(t)dt$$

The inverse Fourier transform is defined by:

$$\mathcal{F}^{-1}g(t) := \int_{-\infty}^{+\infty} e^{2\pi i st} g(s)ds$$

The Fourier inversion theorem states that

$$\mathcal{F}(\mathcal{F}^{-1}g) = g, \quad \mathcal{F}^{-1}(\mathcal{F}f) = f$$

Riemann Lebesgue Theorem If $f \in \mathcal{L}^1(\mathbb{R})$ then

$$\lim_{|s| \rightarrow \infty} \mathcal{F}(s) = 0$$

Riemann Lebesgue Theorem If $f \in \mathcal{L}^1(\mathbb{R})$ then

$$\lim_{|s| \rightarrow \infty} \mathcal{F}(s) = 0$$

Plancherel Theorem If $f \in \mathcal{L}^1(\mathbb{R}) \cap \mathcal{L}^2(\mathbb{R})$ then $\hat{f} \in \mathcal{L}^2(\mathbb{R})$ and

$$\int_{-\infty}^{\infty} |f(t)|^2 dt = \int_{-\infty}^{\infty} |\hat{f}(s)|^2 ds$$

Riemann Lebesgue Theorem If $f \in \mathcal{L}^1(\mathbb{R})$ then

$$\lim_{|s| \rightarrow \infty} \mathcal{F}(s) = 0$$

Plancherel Theorem If $f \in \mathcal{L}^1(\mathbb{R}) \cap \mathcal{L}^2(\mathbb{R})$ then $\hat{f} \in \mathcal{L}^2(\mathbb{R})$ and

$$\int_{-\infty}^{\infty} |f(t)|^2 dt = \int_{-\infty}^{\infty} |\hat{f}(s)|^2 ds$$

Theorem If $f, g \in \mathcal{L}^1(\mathbb{R})$ then

$$\int_{-\infty}^{\infty} \hat{f}(s)g(s)ds = \int_{-\infty}^{\infty} f(x)\hat{g}(x)dx$$

Example 1: Odd and Even functions.

For any even function $f(t) = f(-t)$ the following relation holds

$$\mathcal{F}f(s) = \int_{-\infty}^{+\infty} \cos(2\pi st) f(t) dt = 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt$$

Example 1: Odd and Even functions.

For any even function $f(t) = f(-t)$ the following relation holds

$$\mathcal{F}f(s) = \int_{-\infty}^{+\infty} \cos(2\pi st) f(t) dt = 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt$$

If $f(t) = -f(-t)$ is odd, we have

$$\mathcal{F}f(s) = -i \int_{-\infty}^{+\infty} \sin(2\pi st) f(t) dt = -2i \int_0^{+\infty} \sin(2\pi st) f(t) dt$$

Example 2: Exponential even function Let $f(t) = e^{-|t|}$ then

$$\begin{aligned}\mathcal{F}f(s) &= 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt \\ &= 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt\end{aligned}$$

Example 2: Exponential even function Let $f(t) = e^{-|t|}$ then

$$\begin{aligned}\mathcal{F}f(s) &= 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt \\ &= 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt\end{aligned}$$

Now using Euler formulae

$$\cos \alpha = \frac{e^{i\alpha} + e^{-i\alpha}}{2} \quad \sin \alpha = \frac{e^{i\alpha} - e^{-i\alpha}}{2i}$$

Example 2: Exponential even function Let $f(t) = e^{-|t|}$ then

$$\begin{aligned}\mathcal{F}f(s) &= 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt \\ &= 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt\end{aligned}$$

Now using Euler formulae

$$\cos \alpha = \frac{e^{i\alpha} + e^{-i\alpha}}{2} \quad \sin \alpha = \frac{e^{i\alpha} - e^{-i\alpha}}{2i}$$

we find

$$\mathcal{F}f(s) = 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt = 2 \int_0^{+\infty} e^{-t} \frac{e^{i(2\pi st)} + e^{-i(2\pi st)}}{2} dt$$

$$\mathcal{F}f(s) = \int_0^{+\infty} \left(e^{(-1+2i\pi s)t} + e^{(-1-2i\pi s)t} \right) dt$$

$$\begin{aligned}
 \mathcal{F}f(s) &= \int_0^{+\infty} \left(e^{(-1+2i\pi s)t} + e^{(-1-2i\pi s)t} \right) dt \\
 &= \left[\frac{e^{(-1+2i\pi s)t}}{-1 + 2i\pi s} \right]_{t=0}^{t=\infty} + \left[\frac{e^{(-1-2i\pi s)t}}{-1 - 2i\pi s} \right]_{t=0}^{t=\infty}
 \end{aligned}$$

$$\begin{aligned}
 \mathcal{F}f(s) &= \int_0^{+\infty} \left(e^{(-1+2i\pi s)t} + e^{(-1-2i\pi s)t} \right) dt \\
 &= \left[\frac{e^{(-1+2i\pi s)t}}{-1+2i\pi s} \right]_{t=0}^{t=\infty} + \left[\frac{e^{(-1-2i\pi s)t}}{-1-2i\pi s} \right]_{t=0}^{t=\infty} \\
 &= \frac{1}{1-2i\pi s} + \frac{1}{1+2i\pi s}
 \end{aligned}$$

$$\begin{aligned}
\mathcal{F}f(s) &= \int_0^{+\infty} \left(e^{(-1+2i\pi s)t} + e^{(-1-2i\pi s)t} \right) dt \\
&= \left[\frac{e^{(-1+2i\pi s)t}}{-1 + 2i\pi s} \right]_{t=0}^{t=\infty} + \left[\frac{e^{(-1-2i\pi s)t}}{-1 - 2i\pi s} \right]_{t=0}^{t=\infty} \\
&= \frac{1}{1 - 2i\pi s} + \frac{1}{1 + 2i\pi s} \\
&= \frac{1 + 2i\pi s + 1 - 2i\pi s}{1^2 - (2i\pi s)^2}
\end{aligned}$$

$$\begin{aligned}
\mathcal{F}f(s) &= \int_0^{+\infty} \left(e^{(-1+2i\pi s)t} + e^{(-1-2i\pi s)t} \right) dt \\
&= \left[\frac{e^{(-1+2i\pi s)t}}{-1 + 2i\pi s} \right]_{t=0}^{t=\infty} + \left[\frac{e^{(-1-2i\pi s)t}}{-1 - 2i\pi s} \right]_{t=0}^{t=\infty} \\
&= \frac{1}{1 - 2i\pi s} + \frac{1}{1 + 2i\pi s} \\
&= \frac{1 + 2i\pi s + 1 - 2i\pi s}{1^2 - (2i\pi s)^2} \\
&= \frac{2}{1 + 4\pi^2 s^2}
\end{aligned}$$

Example 2: Exponential even function Let $f(t) = e^{-|t|}$, then

$$\begin{aligned}\mathcal{F}f(s) &= 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt \\ &= 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt\end{aligned}$$

Example 2: Exponential even function Let $f(t) = e^{-|t|}$, then

$$\begin{aligned}\mathcal{F}f(s) &= 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt \\ &= 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt \\ &= \left[\frac{2e^{-t}(2\pi s \sin(2\pi st) - \cos(2\pi st))}{4\pi^2 s^2 + 1} \right]_{t=0}^{t=+\infty}\end{aligned}$$

Example 2: Exponential even function Let $f(t) = e^{-|t|}$, then

$$\begin{aligned}\mathcal{F}f(s) &= 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt \\ &= 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt \\ &= \left[\frac{2e^{-t}(2\pi s \sin(2\pi st) - \cos(2\pi st))}{4\pi^2 s^2 + 1} \right]_{t=0}^{t=+\infty}\end{aligned}$$

since $\int e^{-t} \cos(At) dt = \frac{e^{-t}(A \sin At - \cos At)}{1 + A^2}$

Example 2: Exponential even function Let $f(t) = e^{-|t|}$, then

$$\begin{aligned}\mathcal{F}f(s) &= 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt \\ &= 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt \\ &= \left[\frac{2e^{-t}(2\pi s \sin(2\pi st) - \cos(2\pi st))}{4\pi^2 s^2 + 1} \right]_{t=0}^{t=+\infty}\end{aligned}$$

since $\int e^{-t} \cos(At) dt = \frac{e^{-t}(A \sin At - \cos At)}{1 + A^2}$ so that

$$\mathcal{F}f(s) = \frac{2}{1 + 4\pi^2 s^2}$$

Example 2: Exponential even function Let $f(t) = e^{-|t|}$, then

$$\begin{aligned}\mathcal{F}f(s) &= 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt \\ &= 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt\end{aligned}$$

Example 2: Exponential even function Let $f(t) = e^{-|t|}$, then

$$\begin{aligned}\mathcal{F}f(s) &= 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt \\ &= 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt \\ &= \left[\frac{2e^{-t}(2\pi s \sin(2\pi st) - \cos(2\pi st))}{4\pi^2 s^2 + 1} \right]_{t=0}^{t=+\infty}\end{aligned}$$

Example 2: Exponential even function Let $f(t) = e^{-|t|}$, then

$$\begin{aligned}\mathcal{F}f(s) &= 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt \\ &= 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt \\ &= \left[\frac{2e^{-t}(2\pi s \sin(2\pi st) - \cos(2\pi st))}{4\pi^2 s^2 + 1} \right]_{t=0}^{t=+\infty}\end{aligned}$$

since $\int e^{-t} \cos(At) dt = \frac{e^{-t}(A \sin At - \cos At)}{1 + A^2}$

Example 2: Exponential even function Let $f(t) = e^{-|t|}$, then

$$\begin{aligned}\mathcal{F}f(s) &= 2 \int_0^{+\infty} \cos(2\pi st) f(t) dt \\ &= 2 \int_0^{+\infty} e^{-t} \cos(2\pi st) dt \\ &= \left[\frac{2e^{-t}(2\pi s \sin(2\pi st) - \cos(2\pi st))}{4\pi^2 s^2 + 1} \right]_{t=0}^{t=+\infty}\end{aligned}$$

since $\int e^{-t} \cos(At) dt = \frac{e^{-t}(A \sin At - \cos At)}{1 + A^2}$ so that

$$\mathcal{F}f(s) = \frac{2}{1 + 4\pi^2 s^2}$$